



## Bottom-up effects of lake sediment on pelagic food-web compartments: a mesocosm study

Loïc Harrault, Béatrice Allard, Jacques Mériguet, David Carmignac, Sylvain Huon, Benoît Gauzens, Gérard Lacroix

### ► To cite this version:

Loïc Harrault, Béatrice Allard, Jacques Mériguet, David Carmignac, Sylvain Huon, et al.. Bottom-up effects of lake sediment on pelagic food-web compartments: a mesocosm study. *Freshwater Biology*, 2014, 59 (8), pp.1695-1709. 10.1111/fwb.12375 . hal-01068352

**HAL Id: hal-01068352**

**<https://hal.science/hal-01068352>**

Submitted on 20 Nov 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Bottom-up effects of lake sediment on pelagic compartments: a mesocosm study**

Journal:	<i>Freshwater Biology</i>
Manuscript ID:	FWB-P-Apr-13-0212.R2
Manuscript Type:	Standard Paper
Date Submitted by the Author:	09-Mar-2014
Complete List of Authors:	Harrault, Loïc; CNRS, allard, beatrice; cnrs, Mériguet, Jacques; ENS, Carmignac, David; ENS, Huon, Sylvain; UPMC, Gauzens, Benoit; ENS, Iacroux, gerard
Keywords:	Lakes < Habitat, Population < Level of Organisation, Molecular < Level of Organisation, Ecosystem < Level of Organisation, Biomanipulation < Applied Issues, Eutrophication < Applied Issues, Biochemical analysis < Process / Approach / Methods, Food webs < Process / Approach / Methods

SCHOLARONE™  
Manuscripts

1  
2  
3 1 **Bottom-up effects of lake sediment on pelagic compartments: a mesocosm study**  
4  
5

6 2  
7

8  
9 3 Loïc Harrault<sup>1,2,3,#</sup>, Béatrice Allard<sup>1</sup>, Jacques Mériguet<sup>3</sup>, David Carmignac<sup>3</sup>, Sylvain Huon<sup>1</sup>,  
10  
11 4 Benoit Gauzens<sup>3,4</sup> and Gérard Lacroix<sup>5,6</sup>  
12  
13

14 <sup>1</sup> Université Pierre et Marie Curie, UMR CNRS 7618 - Bioemco, 4 place Jussieu, 75252 Paris  
15  
16 Cedex05, France  
17  
18

19  
20 <sup>2</sup> Université Rennes 1, UMR CNRS 6118 - Géosciences Rennes, Campus de Beaulieu, 35042  
21  
22 Rennes Cedex, France  
23  
24

25 <sup>3</sup> Ecole Normale Supérieure, UMR CNRS 7618 - Bioemco, 46 rue d’Ulm, 75005 Paris,  
26  
27 France  
28  
29

30 <sup>4</sup> Université Rennes 1, UMR CNRS 6553 ECOBIO, Campus de Beaulieu, 35042 Rennes  
31  
32 Cedex, France  
33  
34

35 <sup>5</sup> CNRS, UMR iEES - Paris, Institute of ecology and environmental sciences - Paris,  
36  
37 Université Pierre et Marie Curie, 7 quai St.-Bernard, 75005 Paris, France  
38  
39

40 <sup>6</sup> Ecole Normale Supérieure, UMS CNRS 3194 - CEREAP – Ecotron IleDeFrance, 78 rue du  
41  
42 Château, 77140 St-Pierre-lès-Nemours, France  
43  
44  
45

46  
47  
48 □ corresponding author : loic.harrault@gmail.com  
49

50 **Keywords:** sediment biodegradability, organic matter, lipid biomarkers, stoichiometry,  
51  
52 nutrient release, mesocosms, pelagic compartments  
53  
54

55  
56  
57 **Abbreviated title:** Influence of lake sediment on pelagic compartments  
58  
59  
60

## Summary

1. Sediment plays a key role in organic matter (OM) and internal nutrient cycling in lakes. The role of sediment as a source of OM, and its potential bottom-up effects on the pelagic food web have been rarely studied. Particularly, the influence of the biochemical composition of sediment OM on pelagic compartments remains largely unknown.

2. In a five-months experiment, we studied the influence of two different sediments added at the bottom of large replicated mesocosms on the biomass, the elemental and the lipid compositions of seston and zooplankton. The influence of sediment treatments on sedimentation rates, elemental and biochemical compositions and potential biodegradability of recently sedimented OM (ca. 1 week) was also examined. The two added sediments ( $S_1$  and  $S_2$ ) presented very contrasted elemental and biochemical compositions and potential biodegradabilities. According to their contents in organic carbon, nitrogen, proteins, sugars and polyunsaturated fatty acids,  $S_2$  appeared to be much more biodegradable than  $S_1$ . Therefore, the  $S_2$  sediment was expected to release more nutrients and OM to the water column than  $S_1$ , leading to changes in communities, stoichiometry and lipid compositions of pelagic compartments.

3. Probably due to its very poor content in labile compounds, the presence of  $S_1$  at the bottom of the mesocosms did not induce changes in the biomass of seston and zooplankton. Only few changes in the stoichiometry of these compartments were observed. On the contrary,  $S_2$  sediment released more phosphorus and dissolved OM in the water column than  $S_1$ . As a result,  $S_2$  treatment induced an increase in seston biomass and therefore, in zooplankton biomass *via* herbivory.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 47 4. None of the sediment treatments affected the lipid composition of seston and
- 48 zooplankton. Moreover, neither S<sub>1</sub> nor S<sub>2</sub> induced changes in the sedimentation rates,
- 49 elemental and lipid compositions, and potential biodegradability of recent sediments.
- 50 Our mesocosm experiment suggests that differences in the quality of lake sediment
- 51 lead to moderate changes in the pelagic communities in the absence of planktivorous
- 52 or omnivorous fish.
- 53 5. Our results might explain the efficiency of biomanipulations for improving water
- 54 quality of eutrophic lakes despite potential nutrient release from sediment. Finally, our
- 55 results provide additional support for the ecological significance of mesocosms for
- 56 studying processes occurring at larger scales.

## 58 Introduction

59 Since a few decades, the role of sediment in the biological and biogeochemical processes in  
60 lacustrine ecosystems has received an increased attention. Lipid composition of sedimented  
61 organic matter (SOM) has been widely used to assess, via specific biomarkers, i)  
62 autochthonous vs allochthonous origin of sediment (Canuel & Martens, 1993; Bechtel &  
63 Schubert, 2009; Castaneda & Schouten, 2011), ii) transfers of energy and nutrients between  
64 primary producers, herbivores and other consumers within aquatic food webs (Müller-  
65 Navarra *et al.*, 2000; Masclaux *et al.*, 2009) or iii) inter-specific differences (Volkman *et al.*,  
66 1988; Cranwell *et al.*, 1990). The sediment geochemistry has also been used to study the  
67 carbon balance in lacustrine ecosystems and its contribution to the global carbon cycle (Alin  
68 & Johnson, 2007; Cole *et al.*, 2007). The net heterotrophy of lakes (CO<sub>2</sub> source for  
69 atmosphere) is the most commonly observed carbon fate in concerned studies (del Giorgio *et al.*,  
70 1997; Cole *et al.*, 2000). It has been shown that this balance can be influenced by food-  
71 web structure, nutrient concentration (Schindler *et al.*, 1997), or climate (Kosten *et al.*, 2010),  
72 which, in turn, modifies sedimentation (Flanagan *et al.*, 2006). Furthermore, several recent  
73 studies in both marine mesocosms (Canuel *et al.*, 2007; Spivak *et al.*, 2007) and freshwater  
74 mesocosms (Allard *et al.*, 2011; Danger *et al.*, 2012), have shown that the biochemical  
75 composition (e.g. organic carbon, protein, sugar and lipid contents) of recently deposited  
76 sediments depends on food-web structure.

77 Once sedimented, OM is subject to transformation and biodegradation by benthic  
78 macroinvertebrates (Mermillod-Blondin *et al.*, 2003; Nogaro *et al.*, 2008) and  
79 microorganisms (Gächter *et al.*, 1988; Hargreaves, 1998). Biodegradation depends on several  
80 parameters, such as dissolved oxygen concentration at the sediment-water interface, binding  
81 on mineral matrix (Wakeham & Canuel, 2006), physical resuspension (Sanford, 1992), or OM  
82 composition (Hulthe *et al.*, 1998). This latter has been largely used to provide information on

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

83 the degradation state and the biodegradability of the sediment, due to the large range of  
84 lability covered by the different components of OM (Meckler *et al.*, 2004; Schubert *et al.*,  
85 2005). During its degradation, the sediment can release into the water column nutrients and  
86 dissolved organic matter (DOM; Reynolds, 1996; Klump *et al.*, 2009), which will support  
87 both primary and bacterial productions (del Giorgio & Cole, 1998). As phytoplankton and  
88 bacteria are the basal resources for aquatic food webs, release of nutrients and OM from  
89 sediment might strongly impact food-web communities via a bottom-up forcing. Whereas the  
90 bottom-up effects of inorganic nutrients or light on aquatic ecosystems are well known (see  
91 for example, Urabe *et al.*, 2002; Cebrian & Lartigue, 2004; Hessen *et al.*, 2004; Dickman *et*  
92 *al.*, 2006; Spivak *et al.*, 2007), those of SOM have been more rarely studied. In natural lakes,  
93 it has been shown that the temporal dynamics of bacterioplankton communities are strongly  
94 dependent on the DOM origin and amount (Berdjeb *et al.*, 2011), suggesting a bottom-up  
95 forcing from DOM. As DOM can originate from sediment biodegradation, one may wonder if  
96 sediment could be responsible for a bottom-up control for the whole food web, through the  
97 availability of resources for basal organisms (e.g. phytoplankton and bacterioplankton).  
98 Moreover, as the sediment biodegradability seems to be linked to its OM composition (Allard  
99 *et al.*, 2011; Danger *et al.*, 2012; Harrault *et al.*, 2012), an interesting issue is whether  
100 different sediment compositions could lead to different bottom-up forcing on aquatic food  
101 webs.

102 In this paper, we report results of a lake mesocosm study examining how the presence  
103 and the OM composition of sediment can exert a bottom-up effect on the composition of  
104 seston, zooplankton and recently deposited (ca. 1 week) sediment. Three treatments were  
105 compared: enclosures without added sediment ( $S_0$ ), enclosures with carbon-poor sediment  
106 ( $S_1$ ), and enclosures with carbon-rich sediment ( $S_2$ ). Sediment  $S_1$  was taken in the bottom of  
107 Lake Créteil (Lacroix & Lescher-Moutoué, 1995), while  $S_2$  came from a previous mesocosm

experiment (Danger *et al.*, 2008, 2012) and was composed of OM from pelagic origin, settled down during a few years in large pelagic enclosures. The compounds that are preferentially degraded by bacteria, such as polyunsaturated fatty acids (PUFAs, Cranwell, 1981), and sugars and proteins (Weiss & Simon, 1999), were much more abundant in S<sub>2</sub> than in S<sub>1</sub> (Table 1). This strongly suggested that S<sub>2</sub> was much more biodegradable than S<sub>1</sub>. Indeed, S<sub>1</sub> had been formed over several decades in a shallow gravel-pit lake with a clear tendency to oligotrophication since the end of its exploitation (Garnier, 1992). In contrast, the composition of S<sub>2</sub> was typical of sediments found in hypereutrophic conditions (Søndergaard, 1988), in aquaculture areas (Yokoyama *et al.*, 2009), and in extensive fish ponds (Banas *et al.*, 2008).

During 5 months, the biomass and the elemental composition of pelagic compartments were determined. At the end of the experiment, the lipid composition of the biotic compartments was determined. According to the results of Spivak *et al.* (2007), Allard *et al.* (2011), Danger *et al.* (2012) and Harrault *et al.* (2012), we hypothesised that:

- (i) a positive bottom-up forcing from sediment would increase the biomass of seston (direct effect through the ingestion of released OM and/or nutrients) and the biomass of zooplankton (indirect effect through the ingestion of increased seston biomass);
- (ii) the strength of this bottom-up effect would depend on the OM composition of the bottom sediment:  $S_0 < S_1 < S_2$  (i.e. the more biodegradable a sediment, the greater the bottom-up forcing);
- (iii) this effect could change the stoichiometry and/or the lipid biomarker composition of pelagic compartments by modifying nutrient balances.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Methods**

Study site, experimental design and choice of bottom sediments

This study took place in Lake Créteil (48°46'37''N, 2°26'47''E), a small (42 ha), shallow (mean depth 4 m) sandpit lake 15 km southeast of Paris, France (for more information on this lake, see Danger *et al.*, 2008). Twenty-four translucent polyethylene enclosures, sealed at the bottom, were suspended above the lake surface on a floating pontoon. The volume of each enclosure was ca. 13.5 m<sup>3</sup> (2 × 1.5 × 4.5 m depth). At the end of November 2009, enclosures were randomly filled in successive steps with pumped lake water. To study the bottom-up effects of initial sediment on pelagic compartments and on recently deposited sediment, two contrasted sediments were set down at the bottom of the enclosures. Enclosures without added sediment were used as a control (S<sub>0</sub>). Each treatment was replicated eight times. The first sediment (S<sub>1</sub>) was collected randomly in Lake Créteil with an Uwitec bottom sampler and consisted of a dark grey sandy material. The second sediment (S<sub>2</sub>) was collected in enclosures used in a previous study (Danger *et al.*, 2008, 2012). S<sub>1</sub> and S<sub>2</sub> were chosen to simulate contrasted bottom-up treatments in enclosures. Both autochthonous and allochthonous sources contributed to S<sub>1</sub>. This sediment had been accumulated and exposed to microbial and benthic reworking over several decades. In contrast, S<sub>2</sub> was sampled in enclosures sealed at their bottom and largely preserved from allochthonous inputs, and therefore mainly of autochthonous origin. Moreover, this sediment is relatively young since it was collected in experimental mesocosms set up in 2005 (Danger *et al.*, 2008) and has been exposed to degradation over a shorter period than the sediment of the lake. The difference in the initial sediment quality was estimated using the sugar, protein and PUFA contents (Table

1  
2  
3 155 1). The difference in the degradation state of the initial sediments was estimated using their  
4  
5 156 bacterial fatty acid (BACTFA),  $\alpha$ - and  $\beta$ -hydroxy fatty acid (OH-FA) and stanol contents  
6  
7 157 (Supplementary Table 1).  
8  
9

10 158 Once collected, sediments (ca. 600 L each) were pooled into 200-L tanks for homogenisation.  
11  
12 159 Trays ( $2 \times 1.5$  m), filled with ca. 70 L of homogenized sediment (ca. 2.4 cm depth layer),  
13  
14 160 were lowered very slowly at the bottom of each enclosure to minimize sediment resuspension  
15  
16 161 and subsequent dissolution of nutrients and OM. Empty trays were put in the control  
17  
18 162 enclosures. To allow sedimentation of suspended material in  $S_1$  and  $S_2$  enclosures, we started  
19  
20 163 samplings on February 2010.  
21  
22  
23  
24  
25  
26

27 165 Orthophosphates, ammonium and nitrates analyses  
28  
29  
30  
31 166

32  
33 167 Orthophosphate, ammonium and nitrate concentrations were determined for water sampled in  
34  
35 168 May and June 2010 using the phosphorus-ammonium molybdate and the indephenol blue  
36  
37 169 spectrometric methods, respectively (AFNOR, 1990, 2004).  
38  
39  
40  
41  
42  
43

44 171 Dissolved organic carbon analysis  
45  
46  
47 172

48  
49  
50 173 Water samples collected in May 2010 were filtered through a Whatman GF/F filter. Dissolved  
51  
52 174 organic carbon (DOC) concentration was determined using a total organic carbon analyser  
53  
54 175 (TOC-5000A, Shimadzu, Kyoto, Japan).  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

177 Seston and zooplankton sampling

178

179 Seston biomass and elemental composition were determined monthly. Water was sampled

180 monthly from February to June 2010 at different depths and locations with a 2-L sampling

181 bottle (Uwitec) in each enclosure. Water samples were filtered through a 50-µm nylon filter to

182 remove zooplankton, and then filtered through a pre-weighted Whatman GF/F glass-fiber

183 filter (nominal cut-off: 0.7 µm) in order to gather seston (particulate matter between 0.7 and

184 50 µm). Filters were dried overnight at 60°C and weighted to determine seston biomass. Dry

185 filters were stored in the dark at room temperature until elemental and lipid analyses.

186 Zooplankton biomass was determined monthly by sampling 60 L of water at different depths

187 and locations in each enclosure with a 12-L Schindler plankton trap equipped with a 50-µm

188 filter. Zooplankton was gathered in GF/F-filtered water for several hours to allow evacuation

189 of gut content. Zooplankton was concentrated on a 50-µm filter, washed with deionized water

190 to remove particles and dissolved matter bound to their shell, placed on a pre-weighted

191 Whatman GF/A glass-fiber filter (nominal cut-off: 1.6 µm), and dried overnight at 60°C. Dry

192 zooplankton was ground and stored in the dark at room temperature until elemental and lipid

193 analyses.

194

195 Zooplankton counting

196

197 Zooplankton composition was determined in February and May 2010 by filtering 60 L of

198 water at different depths and locations in three enclosures of each treatment with a 12-L

199 Schindler plankton trap equipped with a 50-µm filter. Zooplankton was preserved in 4%

formaldehyde. Zooplankton taxa were identified and counted under a Leica stereo-microscope on subsamples at different dilutions in Dollfuss chambers. Copepods were separated into cyclopidae, calanoida and nauplii. Cladocera were separated into *Daphnia*, *Ceriodaphnia*, and Chydoridae. Rotifers were counted globally.

204

205 Sampling of recent sediment

206

Recently deposited sediments were sampled in May and June 2010. Sedimentation rates were determined using six sediment traps deployed in each enclosure. Traps consisted of 5-cm diameter and 30-cm long PVC tubes, suspended at 4 m depth. Suspended 0.5 m above the bottom of the enclosures, sediment traps were preserved for bottom sediment resuspension. Traps were deployed for 7- to 9-day intervals. Material collected from the six traps was pooled in a collection flask and allowed to sediment overnight at 4°C. The supernatant and zooplankton therein was removed. Sediment was freeze-dried, weighted, ground and stored in the dark at room temperature until analyses. Sedimentation rates were calculated as the mass of dry matter divided by the duration of the trap deployment and the total surface of the six traps, and expressed in  $\text{g m}^{-2} \text{ day}^{-1}$ .

217

218 Elemental composition of seston, zooplankton, initial and recent sediment

219

Carbon and nitrogen contents of dried samples were determined using a CHN elementary analyzer (FlashEA 1112 series, Thermo Fisher Scientific) with acetanilide as standard. Initial

1  
2  
3 222 and recent sediment organic carbon contents (OC) were determined after removal of inorganic  
4  
5 223 carbon (IC) from sediment by successive additions of 1 M HCl (Hedges & Stern, 1984).  
6  
7  
8 224  
9  
10  
11 225 Sugar and protein colorimetric assays of sediments  
12  
13  
14 226  
15  
16  
17 227 Freeze-dried sediments were extracted with H<sub>2</sub>O at 100°C for 2 h. The mixture was filtered  
18  
19 228 through a Whatman GF/F filter, and the filtrate was freeze-dried. The freeze-dried aqueous  
20  
21 229 extract was dissolved in a known volume of H<sub>2</sub>O and assayed for sugars and proteins. Sugar  
22  
23 230 contents were determined using the phenol–sulfuric acid colorimetric method with glucose as  
24  
25 231 standard (Dubois *et al.*, 1956). The protein contents were determined using the colorimetric  
26  
27 232 method of Lowry with bovine serum albumin as standard (Lowry *et al.*, 1951). Sugar and  
28  
29 233 protein contents were expressed as percentages of the sediment dry weight (DW).  
30  
31  
32  
33 234  
34  
35 235 Lipid analysis  
36  
37  
38 236  
39  
40  
41  
42 237 All chemicals used were of analytical grade.  
43  
44  
45 238       Apart from initial sediments S<sub>1</sub> and S<sub>2</sub>, analyses of lipids were carried out on three  
46  
47 239 replicates of each treatment for seston, zooplankton and recent sediment sampled in May  
48  
49 240 2010.  
50  
51  
52 241       The filter with collected seston was extracted with a dichloromethane  
53  
54 242 (DCM)/methanol (MeOH) (2/1, v/v) mixture at room temperature for 18 h. The mixture was  
55  
56 243 filtered through a Whatman GF/F glass-fiber filter and solvent was removed under reduced  
57  
58  
59  
60

pressure. Extracts were saponified at 80°C for 2 h with 1 M KOH in MeOH. The pH of saponified extract was brought to 2 by addition of 6 M HCl and lipids extracted with DCM. The organic phase was washed with deionized water until neutral pH, dried over Na<sub>2</sub>SO<sub>4</sub> and DCM removed under reduced pressure. Saponified lipids were treated with ca. 4 M HCl in MeOH at 80°C for 1 h to convert carboxyl groups into their methyl ester derivatives. Esterified extract was then treated with a mixture of anhydrous pyridine/N,O-Bis(trimethylsilyl)trifluoroacetamide (BSTFA) (10/1, v/v) for 10 min at 60°C to convert hydroxyl groups into trimethylsilyl (TMS) ether groups. Lipid components (as methyl esters and TMS ethers) were analysed by gas chromatography-mass spectrometry (GC–MS) with an Agilent 6890 gas chromatograph coupled to an Agilent 5973N mass spectrometer with electron ionization at 70 eV. Separation was achieved using a fused silica column coated with RTX5SilMS (30 m, i.d. 0.25 mm, film thickness 0.5 µm) with helium as carrier gas. The GC oven was programmed from 100 to 320°C at 4°C min<sup>-1</sup>. Unsaturated fatty acids were analysed using gas chromatography-flame ionization detector (GC-FID) and identified by comparison with standards (Larodan, Malmö, Sweden) using fused silica column coated with DB23 (60 m, i.d. 0.25 mm, film thickness 0.25 µm) with helium as carrier gas. The GC oven was programmed from 100 to 170°C at 6.5°C min<sup>-1</sup>, then to 215°C at 2.75°C min<sup>-1</sup> and to 230°C at 40°C min<sup>-1</sup>. Individual component contributions were determined by comparison of the peak areas from GC–MS traces. This method allowed a comparison of the relative abundance of each compound between the different samples analysed, but did not allow the quantification of individual compounds within the total lipids. Dried zooplankton samples were extracted and lipids analyzed as described for seston. The amounts of extracted lipids were too low to allow an accurate weight determination.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Freeze-dried sugar- and protein-free sediment samples were extracted with a chloroform/MeOH (2/1, v/v) mixture for 2 h at 80°C. Lipid extracts were analysed as described for seston.

Statistical analyses

Statistical analyses were performed using R software ([www.r-project.org](http://www.r-project.org)). A linear mixed-effects model (LME) with sediment treatment, time and cross-effect as explanatory variables and replicates as a random effect was used to test the individual and the combined effects of sediment treatment (S<sub>0</sub>, S<sub>1</sub> and S<sub>2</sub>) and time on the biomass of seston and zooplankton (from February to June), zooplankton composition (February and May), sedimentation rates (May and June), elemental compositions of seston, zooplankton (from February to June) and recent sediment (May and June), and orthophosphate, ammonium and nitrate concentrations of water (May and June). For response variables analysed only in May (DOC and lipid biomarkers), effects of sediment treatment were tested with a LME with sediment treatment as explanatory variable and replicates as a random effect. Post-hoc Tukey's tests (Tukey) were performed to compare one treatment to another. Data were log-transformed when necessary to homogenize distributions and variances.

Technical constraints

The design of this experiment had been initially planned to last for almost 2 years and not for 5 months as presented here. With the design initially conceived, regular sampling on various

290 biotic and abiotic parameters were planned from May 2010 to December 2011. Unfortunately,  
291 extreme climatic events have damaged the enclosures, forcing us to stop the experiment on  
292 July 2010 with an incomplete and heterogeneous set of data. As a consequence, phosphorus  
293 analyses in the various compartments of the mesocosms were not possible, and other analyses  
294 (nutrients, DOC, particles, zooplankton taxonomy) were conducted only on one or two  
295 sampling dates.

296

## 297 **Results**

298

299 Water: nutrient and DOC concentrations

300

301 In May and June, sediment treatment had a significant effect on orthophosphate concentration  
302 of water (Table 2). Orthophosphate concentration from S<sub>2</sub> enclosures was significantly higher  
303 than those from S<sub>0</sub> and S<sub>1</sub> enclosures, which did not differ significantly each other.

304 Sediment treatment had no significant effect on the concentrations of ammonium and  
305 nitrates and on the inorganic nitrogen (sum of ammonium and nitrate  
306 concentrations)/orthophosphates ratio (IN/P, Table 2). Despite this absence of significant  
307 effect, a IN/P ratio from S<sub>2</sub> enclosures (5) much lower than that from both S<sub>0</sub> and S<sub>1</sub> (16 and  
308 18, respectively) was observed.

309 In May, sediment treatment had a significant effect on DOC concentration (Table 2).  
310 DOC concentration in S<sub>2</sub> enclosures was significantly higher than that in S<sub>1</sub> enclosures. In



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

311 contrast DOC concentration in S<sub>0</sub> and S<sub>1</sub> enclosures did not differ significantly and DOC  
312 concentration in S<sub>2</sub> enclosures did not differ significantly from that in S<sub>0</sub> enclosures.

313  
314 Seston: biomass and elemental composition

315  
316 Biomass and elemental compositions of seston (particulate matter between 0.7 and 50 µm)  
317 from the different treatments were reported in Table 2.

318 Sediment treatment had a significant effect on seston biomass (Fig. 1a) since, on  
319 average, seston biomass in S<sub>0</sub> and S<sub>1</sub> enclosures did not differ significantly but were lower  
320 than those from S<sub>2</sub> enclosures. Seston biomass was higher during winter than during spring.

321 The C content of seston did not differ significantly among treatments and was slightly  
322 higher on April than for the other dates (Fig. 2a). The N content of seston was affected by  
323 sediment treatment: it did not differ between S<sub>0</sub> and S<sub>1</sub> enclosures but was lower in S<sub>0</sub>  
324 enclosures compared to S<sub>2</sub> enclosures, without any time effect (Fig. 2b). The C/N ratio of  
325 seston from S<sub>0</sub> treatment was higher than that of seston from S<sub>1</sub> and S<sub>2</sub> treatments but this  
326 difference was only observed in February and March (Fig. 2c). The C/N ratio of seston was  
327 lower in February than during the following months.

328  
329 Zooplankton: biomass, specific and elemental compositions

330  
331 Biomass and elemental compositions of zooplankton from the different treatments were  
332 reported in Table 2.

1  
2  
3 333 Sediment treatments had a very significant effect on zooplankton biomass (Fig. 1b).  
4  
5 334 On average, zooplankton biomass did not differ significantly between S<sub>0</sub> and S<sub>1</sub> enclosures  
6  
7 335 but was higher in S<sub>2</sub> enclosures than in S<sub>0</sub> and S<sub>1</sub> ones, especially in March and April (Fig.  
8  
9 336 1b). Zooplankton biomass increased from February to April and then decreased up to June.

10  
11  
12 337 Species composition of zooplankton communities was weakly affected by sediment  
13  
14 338 treatments since only the concentration of Chydoridae was significantly higher in the S<sub>2</sub>  
15  
16 339 enclosures than in S<sub>0</sub> and S<sub>1</sub> ones (Table 4). On average, the number of individuals was lower  
17  
18 340 in February than in May, especially for nauplii of copepods ( $7.3 \pm 3.4$  and  $180.9 \pm 77.7$   
19  
20 341 individuals L<sup>-1</sup>, for February and May, respectively. C and N contents and C/N ratio of  
21  
22 342 zooplankton from S<sub>0</sub>, S<sub>1</sub> and S<sub>2</sub> treatments did not differ significantly (Table 2). The C  
23  
24 343 content of zooplankton slightly increased from February to March and then slightly decreased  
25  
26 344 until June (Fig. 3a). The N content of zooplankton sampled in February was lower than that  
27  
28 345 sampled on the other dates (Fig. 3b). The opposite trend was observed for the C/N ratio of  
29  
30 346 zooplankton (Fig. 3c).

31  
32  
33  
34  
35  
36 347

37  
38 348 Recent sediment: sedimentation rates, elemental compositions and sugar and protein contents

39  
40  
41  
42 349

43  
44 350 Recent sediments were sampled in May and in June. Sedimentation rates and elemental  
45  
46 351 composition of recent sediments were reported in Table 2.

47  
48  
49 352 Sediment treatments and time did not significantly affect the sedimentation rates (Fig.  
50  
51 353 1c), although sedimentation rates tended to be higher in S<sub>1</sub> and S<sub>2</sub> than in S<sub>0</sub> enclosures.

52  
53  
54 354 Sediment treatments affected the C, OC and N contents of recent sediments since  
55  
56 355 those from S<sub>1</sub> enclosures were lower than those from S<sub>0</sub> and S<sub>2</sub> ones (Fig. 4a and 4b). The

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

C/N ratio of recent sediment from S<sub>1</sub> treatment was higher than that from S<sub>0</sub> and S<sub>2</sub> treatments (Fig. 4c). In contrast, the OC/N ratio was not significantly affected by the sediment treatment.

The amount of sugar and protein contents of recent sediments did not differ significantly between the different treatments (Table 2).

Lipid compositions of seston, zooplankton and recent sediments

Mean lipid distributions of the three pelagic compartments were reported in Table 3.

Except for chlorophyll-derived compounds in recent sediments (LME:  $p = 0.026$ ,  $F_{2,6} = 7.13$ ), sediment treatments had no significant effect on the relative amounts of the different lipid classes of seston, zooplankton and recent sediments (data not shown). Chlorophyll-derived compounds were more abundant in recent sediments from S<sub>2</sub> ( $4.7 \pm 1.7$  % of total lipids) enclosures than in recent sediments from S<sub>0</sub> ( $1.8 \pm 0.7$  % of total lipids, Tukey:  $p = 0.033$ ) and S<sub>1</sub> ( $2.0 \pm 0.5$  % of total lipids, Tukey:  $p = 0.048$ ) enclosures. FAs largely dominated their lipid distributions, followed by alkanols. Sterols, hydroxy acids and chlorophyll-derived compounds were present in lower relative abundances.

Sediment treatments had no effect on the FA distributions (Supplementary Tables 1 and 2). For all treatments saturated fatty acids (SAFAs) dominated the distribution, followed by monounsaturated fatty acids (MUFAs), PUFAs (long-chain PUFAs, carbon number  $\geq 20$ , accounting for ca. 1, 27 and 8% of total PUFAs for seston, zooplankton and recent sediments, respectively) and BACTFAs.

377 Sterol distributions did not depend on sediment treatment (Table 3). For all treatments,  
378 24-ethyl-cholest-5-enol ( $C_{29}\Delta^5$ ) and cholesterol ( $C_{27}\Delta^5$ ) largely dominated the sterol  
379 distribution of recent sediments, with stanols accounting for less than 2% of total sterols.

380

## 381 Discussion

382 All the criteria used for the selection of the bottom sediments suggest that the potential  
383 biodegradability of  $S_2$  could have been one order of magnitude higher than that of  $S_1$ . The  
384 pool of OM, which represented the total stock of resources potentially used by bacteria,  
385 proteins and sugars, essential for bacterial growth, and PUFAs, preferentially degraded by  
386 bacteria, were clearly more abundant in  $S_2$  than in  $S_1$ .

387

388 Effects of bottom-up forcing on pelagic compartments

389

390 According to the characteristics of  $S_1$  and  $S_2$ , a release of nutrients and OM in the water  
391 column in the order  $S_0 < S_1 < S_2$  was expected. We hypothesized that this would lead to an  
392 enhanced bottom-up forcing on the basal organisms of the food web, resulting in an increase  
393 of seston biomass. By consuming seston, zooplankton was also expected to increase its  
394 biomass with addition of sediment of increasing biodegradability. The phosphate and DOC  
395 concentrations in the water column as well as the biomass values of seston and zooplankton,  
396 all higher in  $S_2$  treatment than in  $S_1$  and  $S_0$  treatments, partially corroborated this hypothesis.

397 The sediment of  $S_2$  enclosures, which was typical of eutrophic to hypertrophic lakes  
398 and fishponds (Søndergaard, 1988; Banas *et al.*, 2008), clearly fostered pelagic compartments.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

399 This result confirms that eutrophic ecosystems are subject to internal nutrient loading, and  
400 that released nutrients may be taken up by aquatic food webs (Søndergaard *et al.*, 2003,  
401 2007). Interestingly, the experiment was run from December to June, when dissolved oxygen  
402 remained oversaturated in all enclosures, even near sediments (data not shown). This  
403 strengthens the statement of Nürnberg (2009) that “phosphorus release is not restricted to and  
404 does not require anoxia in the overlaying water”. As for dissolved oxygen, we did not find  
405 any significant treatment effect on the pH values in the water column, which were high and  
406 ranged between 8.1 and 8.5 (data not shown). Such values are considered to favour  
407 phosphorus release from sediment (Koski-Vahala & Hartikainen, 2001). Other processes may  
408 have favoured nutrient transfer into the overlaying water of S<sub>2</sub> enclosures. Activity of benthic  
409 micro- and macroinvertebrates is probably one of these mechanisms, as suggested by the  
410 higher abundance of Chydoridae in microinvertebrate samples and by the frequent  
411 observation of Chironomids in the S<sub>2</sub> enclosures (direct observations). Moreover, a higher  
412 microbial mineralisation of the highly biodegradable S<sub>2</sub> SOM was expected (Gächter &  
413 Meyer, 1993; Pettersson, 1998). The observed changes in dissolved nutrients in water and  
414 associated biomass of seston suggest a limitation of planktonic primary producers by  
415 phosphorus in S<sub>0</sub> and S<sub>1</sub> enclosures, and a reduction of such P-limitation, accompanied by a  
416 higher primary productivity, in S<sub>2</sub> enclosures. It would be interesting to conduct long-term  
417 experiments in order to verify how internal nutrient loading associated to sediment-water  
418 exchanges may change nutrient limitation of phytoplankton in a seasonal context.

419 By contrast, no difference was observed in the nutrient and DOC concentrations, and  
420 in the biomass of seston and zooplankton between S<sub>1</sub> and S<sub>0</sub> treatments. This absence of  
421 significant difference between S<sub>1</sub> and S<sub>0</sub> suggests that the potential bottom-up effect  
422 associated to OM degradation and nutrient release from the sediment of Lake Créteil was  
423 limited. As highlighted previously (Lacroix *et al.* 1989), the lack of OM accumulation in the

sediment of this shallow sand-pit lake, despite high lake productivity, suggests an efficient OM transfer within the pelagic food web and an efficient nutrient recycling within the water column. The low initial abundance of very labile compounds, such as proteins, sugars and PUFAs in S<sub>1</sub> sediment confirmed this recycling efficiency. Moreover, it is very probable that the natural mineral matrix of Lake Créteil sediment (such mineral matrix was negligible in S<sub>2</sub> enclosures) contained much more P-binding elements, such as iron oxides, which should have favoured a higher retention of phosphorus in the well-oxygenated conditions of the experiment (Mortimer, 1941).

Whatever the treatments, the highest biomass of seston was observed in February and in March, when the biomass and abundance of zooplankton were the lowest. This seasonal trend, previously observed (Danger *et al.*, 2012), agrees with the low grazing activity of zooplankton in late winter, which results in a weak top-down control on phytoplankton. On the other hand, the enhanced grazing activity of zooplankton in spring probably resulted in a more important top-down control on phytoplankton (Sarnelle, 1999).

The bottom-up effect observed for S<sub>2</sub> treatment did not strongly influence the structure of the zooplankton community. The only effect was observed within Cladocera. The typically pelagic daphniidae tended to be less abundant, while the benthic and littoral Chydoridae attained higher densities in the S<sub>2</sub> enclosures. The observed increase in the abundance of Chydoridae in the S<sub>2</sub> enclosures, characterized by their OM-rich sediment, is consistent with the ability of Chydoridae to ingest detritus (de Eyto & Irvine, 2001). Indeed, some Chydoridae, such as *Chydorus* sp., present in the enclosures, are able to use both benthic and open-water food sources (de Eyto & Irvine, 2001). Finally, after a strong increase in the beginning of spring, the biomass of zooplankton from S<sub>2</sub> treatment slowly decreased in the end of spring, suggesting that phytoplankton became rapidly limiting for zooplankton growth in spite of the initial positive bottom-up effect from this sediment. This result supports the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

hypothesis that competition for food might be important for herbivorous zooplankton in such experimental systems when external nutrient loading is absent.

The occurrence of a bottom-up effect of S<sub>2</sub> treatment is also supported by the elemental composition of seston. The higher N content of seston observed in S<sub>2</sub> treatment, compared to S<sub>0</sub> and S<sub>1</sub> ones, suggests a release of N from this “labile” sediment. The lower C/N ratio of seston from S<sub>2</sub> enclosures, probably indicative of a higher nutritional value (Jones *et al.*, 2002), could have contributed to the higher biomass of zooplankton observed in these enclosures. By contrast, zooplankton C/N ratios did not depend on treatments and were quite similar to those obtained by Danger *et al.* (2012). Since species composition of zooplankton was only slightly affected by sediment treatments, this constant C/N ratio is in agreement with the homeostasis constraints on zooplankton stoichiometry. Indeed, zooplankton species have specific elemental compositions (Hessen, 1990; Andersen & Hessen, 1991). Moreover, the decrease in zooplankton C/N ratio with time observed in the enclosures is in agreement with the increase in copepods within the community. As a conclusion, changes in seston stoichiometric ratios do not necessarily induce changes in the structure of the zooplankton community. In a previous study, Danger *et al.* (2012) had shown that changes at the top of the food web affected elemental composition of zooplankton communities by shifting species dominance, but did not affect seston stoichiometric ratios. This suggests that stoichiometric changes are not necessarily transferred up or down along food chains within pelagic food webs.

Although addition of sediment influenced seston stoichiometry, we did not observe any significant influence of the sediment treatments on its lipid composition. On the other hand, previous studies (Allard *et al.*, 2011, Danger *et al.*, 2012) showed that changes in biochemical composition of seston could occur despite similar elemental ratios. This strengthens the conclusion of these authors that both elemental and biochemical compositions

of OM might provide useful information for understanding links between food-web structure and functioning of aquatic ecosystems.

Even if controversies do exist about the importance of internal nutrient loading and the various mechanisms that allow the release of resources from sediments to the overlaying water (see Nürnberg, 2009), our results show that:

(i) sediments with a high abundance of labile OM, which are more typical of eutrophic ecosystems, may clearly foster the pelagic food webs even after the reduction of allochthonous inputs and even in absence of anoxia (Søndergaard *et al.*, 2007);

(ii) sediments with low OM contents and biodegradability, typical of oligo-mesotrophic ecosystems, are necessary for reducing (at least in absence of anoxia) delayed bottom-up forcing of sediment on the pelagic compartments.

Effects of bottom-up forcing on recent sediments

The higher biomass of both seston and zooplankton observed in S<sub>2</sub> enclosures did not induce significant higher sedimentation rates in these enclosures. This might be due to low production of sinking material from seston and zooplankton in pelagic systems characterized by the absence of fish. Indeed, the values obtained in this study are in agreement with those from previous studies carried out in zooplankton-dominated systems (Sarnelle, 1999; Danger *et al.*, 2012). These studies showed that the sedimentation rates are lower in systems dominated by zooplankton than in presence of fish. As a consequence, even if differences in sedimentation rates may have existed, they were probably low and hardly detectable in such fishless and shallow systems.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

497 In our enclosures, the biomass of seston was higher than that of zooplankton.  
498 Moreover, phytoplankton, which constitutes up to 40% of seston (Hessen *et al.*, 2003), is  
499 known to sink much more than zooplankton (Sommer, 1984). Thus, changes in the elemental  
500 composition of seston observed in the different treatments are expected to result in similar  
501 changes in recent sediment. This hypothesis was not supported by the stoichiometric analyses  
502 since the elemental composition of recent sediment from S<sub>2</sub> treatment was similar to that from  
503 S<sub>0</sub> treatment. Surprisingly, the elemental composition of recent sediment from S<sub>1</sub> treatment  
504 differed from that of recent sediment from S<sub>0</sub> and S<sub>2</sub> treatments. At the present time, no  
505 explanation can be put forward for these results.

506 Apart from chlorophyll-derived compounds, no differences in the relative amounts of  
507 lipid classes of recent sediment were observed between the treatments. The higher relative  
508 abundances of chlorophyll-derived compounds in recent sediment from S<sub>2</sub> treatment probably  
509 indicate a higher contribution of phytoplankton to this sediment. The relative abundances of  
510 both FAs and sterols in the recent sediment ranged between those observed for seston and  
511 zooplankton. These results suggest that both seston and zooplankton noticeably contributed to  
512 recent sediment. The relative contribution of each pelagic compartment is difficult to  
513 estimate. As the relative abundances of long-chain PUFAs of recent sediments were quite  
514 low, and more similar to those observed for seston than to those observed for zooplankton,  
515 this suggest a rather low contribution of zooplankton to recent sediment. However, PUFAs  
516 are highly sensitive to bacterial degradation (Cranwell, 1981) and likely do not survive  
517 unaltered for a long time in the water column. Consequently, on the basis of PUFA  
518 concentrations, zooplankton contribution to recent sediment could be underestimated.

519 The low amounts of bacterial FAs suggest a rather low contribution of bacteria to  
520 recent sediments. Nevertheless, stanols were present in higher relative amounts in recent  
521 sediments than in seston and in zooplankton. This indicates that the degradation of settling

organic matter occurred even at the very short time scales (ca. 1 week) of this experiment. However, the relative amounts of unsaturated fatty acids, the most reactive compounds towards biodegradation, were rather high in recent sediments. This suggests that microbial reworking was limited for these very recent sediments.

Lipid biomarkers such as PUFAs and bulk parameters such as sugar and protein contents have been used to estimate and compare the quality of sediments (Canuel *et al.*, 2007; Allard *et al.*, 2011). In the present study, these indicators did not reveal any relationship between the quality of the added sediment and that of the recently deposited one. Finally, the comparison between this study and those dealing with top-down effects (Canuel *et al.*, 2007; Allard *et al.*, 2011; Danger *et al.*, 2012) suggests that the quantity and the quality of settling organic matter in aquatic ecosystems are less affected by bottom-up forces (linked to sediment quality) than by top-down forces (linked to food-web structure) and calls for a factorial approach on these two parameters.

535

## 536 Conclusion

537

This mesocosm experiment demonstrates that the origin and the quality of lake sediment can result in changes in the pelagic compartments. The release of nutrients and OM from the sediment to the water column can induce a bottom-up forcing transferred from seston to zooplankton by means of trophic interactions. However, this effect is substantial only when sediment OM is highly biodegradable and typical of eutrophic systems with high abundances of omnivorous and planktivorous fish (Banas *et al.*, 2008; Yokoyama *et al.*, 2009). This strengthens the suggestion of Harrault *et al.* (2012) that maintaining over a sufficiently long period a top-down control of phytoplankton biomass, for example through the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

biomanipulation of fish communities, should not only induce a more efficient grazing on primary producers, but should also reduce the quantity and biodegradability of SOM, ultimately decreasing internal nutrient cycling. The progressive reduction of internal recycling might explain the acknowledged efficiency of biomanipulations for improving water quality (Jeppesen *et al.*, 2007).

Finally, microcosms and mesocosms have been frequently criticized for their lack of realism (Carpenter, 1996; Schindler, 1998). Our results suggest that the absence of initial sediment in most realized mesocosm experiments did not necessarily induce major biases in the functioning of pelagic systems, and tend to confirm the ecological significance of the effects identified by such manipulations (Spivak & Vanni, 2011). However, it should be kept in mind that other benthic processes, such as redox and pH conditions controlling phosphorus release, or sediment as a source of planktonic organisms and macroinvertebrates, might also strongly impact biogeochemical cycles (Søndergaard *et al.*, 2003). Their importance could be underestimated in most mesocosm experiments.

**Acknowledgments**

This work was supported by the ‘Agence Nationale de la Recherche’ (BIOFUN project ANR-05-BDIV-013-01; PULSE project, ANR-10-CEPL-010). We thank Battle Karimi, Isabelle Godart, Louis Cyrille Gillard, Stéphane Loisel and the team of Lake Créteil for technical support. We thank the anonymous reviewers for their helpful comments. This experiment complies with the current French laws.

**References**

- 569 AFNOR (1990) *Eaux-Méthodes d'Essais*. Association Française de Normalisation, Paris, 736  
570 p.
- 571 AFNOR (2004) Water quality-determination of phosphorus-ammonium molybdate  
572 spectrometric method (EN ISO 6878□:2004).
- 573 Alin S.R. & Johnson T.C. (2007) Carbon cycling in large lakes of the world: A synthesis of  
574 production, burial, and lake-atmosphere exchange estimates. *Global Biogeochemical*  
575 *Cycles*, **21**.
- 576 Allard B., Danger M., Ten-Hage L. & Lacroix G. (2011) Influence of food web structure on  
577 the biochemical composition of seston, zooplankton and recently deposited sediment in  
578 experimental freshwater mesocosms. *Aquatic Sciences*, **73**, 113–126.
- 579 Andersen T. & Hessen D.O. (1991) Carbon, nitrogen, and phosphorus content of freshwater  
580 zooplankton. *Limnology and Oceanography*, **36**, 807–814.
- 581 Banas D., Masson G., Leglise L., Usseglio-Polatera P. & Boyd C.E. (2008) Assessment of  
582 sediment concentration and nutrient loads in effluents drained from extensively managed  
583 fishponds in France. *Environmental pollution*, **152**, 679–85.
- 584 Bechtel A. & Schubert C.J. (2009) A biogeochemical study of sediments from the eutrophic  
585 Lake Lugano and the oligotrophic Lake Brienz, Switzerland. *Organic Geochemistry*, **40**,  
586 1100–1114.
- 587 Berdjeb L., Ghiglione J.F. & Jacquet S. (2011) Bottom-up versus top-down control of hypo-  
588 and epilimnion free-living bacterial community structures in two neighboring freshwater  
589 lakes. *Applied and Environmental Microbiology*, **77**, 3591–3599.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Canuel E.A. & Martens C.S. (1993) Seasonal variations in the sources and alteration of organic matter associated with recently-deposited sediments. *Organic Geochemistry*, **20**, 563–577.

Canuel E.A., Spivak A.C., Waterson E.J. & Duffy J.E. (2007) Biodiversity and food web structure influence short-term accumulation of sediment organic matter in an experimental seagrass system. *Limnology and Oceanography*, **52**, 590–602.

Carpenter S.R. (1996) Microcosm experiments have limited relevance for community and ecosystem ecology. *Ecology*, **77**, 677–680.

Castaneda I.S. & Schouten S. (2011) A review of molecular organic proxies for examining modern and ancient lacustrine environments. *Quaternary Science Reviews*, **30**, 2851–2891.

Cebrian J. & Lartigue J. (2004) Patterns of herbivory and decomposition in aquatic and terrestrial ecosystems. *Ecological Monographs*, **74**, 237–259.

Cole J.J., Pace M. L., Carpenter S. R., Kitchell J. F. (2000) Persistence of net heterotrophy in lakes during nutrient addition and food web manipulations. *Limnology and Oceanography*, **45**, 1718–1730.

Cole J.J., Prairie Y.T., Caraco N.F., McDowell W.H., Tranvik L.J., Striegl R.G., *et al.* (2007) Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget. *Ecosystems*, **10**, 171–184.

Cranwell P.A. (1981) Diagenesis of free and bound lipids in terrestrial detritus deposited in a lacustrine sediment. *Organic Geochemistry*, **3**, 79–89.

- 1  
2  
3 611 Cranwell A., Jaworski G.H.M. & Bickley H.M. (1990) Hydrocarbons, sterols, esters and fatty  
4  
5 612 acids in six freshwater chlorophytes. *Phytochemistry*, **29**, 145–151.  
6  
7  
8  
9 613 Danger M., Lacroix G., Oumarou C., Benest D. & Mériquet J. (2008) Effects of food-web  
10  
11 614 structure on periphyton stoichiometry in eutrophic lakes: a mesocosm study. *Freshwater*  
12  
13 615 *Biology*, **53**, 2089–2100.  
14  
15  
16 616 Danger M., Allard B., Arnous M., Carrias J.-F., Mériquet J., Ten-Hage L., *et al.* (2012)  
17  
18 617 Effects of food-web structure on the quantity and the elemental quality of sedimenting  
19  
20 618 material in shallow lakes. *Hydrobiologia*, **679**, 251–266.  
21  
22  
23  
24 619 Dickman E.M., Vanni M.J. & Horgan M.J. (2006) Interactive effects of light and nutrients on  
25  
26 620 phytoplankton stoichiometry. *Oecologia*, **149**, 676–689.  
27  
28  
29  
30 621 Dubois M., Gilles K.A., Hamilton J.K., Rebers P.A. & Smith F. (1956) Colorimetric method  
31  
32 622 for determination of sugars and related substances. *Analytical Chemistry*, **28**, 350–356.  
33  
34  
35 623 De Eyto E. & Irvine K. (2001) The response of three chydorid species to temperature, pH and  
36  
37 624 food. *Hydrobiologia*, **459**, 165–172.  
38  
39  
40  
41 625 Flanagan K.M., McCauley E. & Wrona F. (2006) Freshwater food webs control carbon  
42  
43 626 dioxide saturation through sedimentation. *Global Change Biology*, **12**, 644–651.  
44  
45  
46 627 Gächter R., Meyer J.S. & Mares A. (1988) Contribution of bacteria to release and fixation of  
47  
48 628 phosphorus in lake sediments. *Limnology and Oceanography*, **33**, 1542–1558.  
49  
50  
51  
52 629 Del Giorgio P.A. & Cole J.J. (1998) Bacterial growth efficiency in natural aquatic systems.  
53  
54 630 *Annual Review of Ecology and Systematics*, **29**, 503–541.  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

631 Del Giorgio, P. A. Cole, J. J., Cimbleris A. (1997) Respiration rates in bacteria exceed  
632 phytoplankton production in unproductive aquatic systems. *Nature*, **385**, 148–151.

633 Garnier J. (1992). Typical and atypical features of phytoplankton in a changing environment -  
634 8 years of oligotrophication in a recently created sand-pit lake (Créteil lake, Paris suburb,  
635 France). *Archiv Fur Hydrobiologie*, **125(4)**, 463–478

636 Goad L.J. & Akihisa T. (1997) *Analysis of Sterols*. Blackie Academic and Professional,  
637 London.

638 Hargreaves J.A. (1998) Nitrogen biogeochemistry of aquaculture ponds. *Aquaculture*, **166**,  
639 181–212.

640 Harrault L., Allard B., Danger M., Maunoury-Danger F., Guilpart A. & Lacroix G. (2012)  
641 Influence of food-web structure on the biodegradability of lake sediment. *Freshwater*  
642 *Biology*, **57**, 2390–2400.

643 Hedges J.I. & Stern J.H. (1984) Carbon and nitrogen determinations of carbonate-containing  
644 solids. *Limnology and Oceanography*, **29**, 657–663.

645 Hessen D.O. (1990) Carbon, nitrogen and phosphorous status in daphnia at varying food  
646 conditions. *Journal of Plankton Research*, **12**, 1239–1249.

647 Hessen D.O., Agren G.I., Anderson T.R., Elser J.J. & De Ruiter P.C. (2004) Carbon,  
648 sequestration in ecosystems: The role of stoichiometry. *Ecology*, **85**, 1179–1192.

649 Hessen D.O., Andersen T., Brettum P. & Faafeng B.A. (2003) Phytoplankton contribution to  
650 sestonic mass and elemental ratios in lakes: Implications for zooplankton nutrition.  
651 *Limnology and Oceanography*, **48**, 1289–1296.

- 652 Hulthe G., Hulth S. & Hall P.O.J. (1998) Effect of oxygen on degradation rate of refractory  
653 and labile organic matter in continental margin sediments. *Geochimica et Cosmochimica*  
654 *Acta*, **62**, 1319–1328.
- 655 Jeppesen E., Meerhoff M., Jacobsen B., Hansen R., Søndergaard M., Jensen J., *et al.* (2007)  
656 Restoration of shallow lakes by nutrient control and biomanipulation-The successful  
657 strategy varies with lake size and climate. *Hydrobiologia*, **581**, 269–285.
- 658 Jones R.H., Flynn K.J. & Anderson T.R. (2002) Effect of food quality on carbon and nitrogen  
659 growth efficiency in the copepod *Acartia tonsa*. *Marine Ecology Progress Series*,  
660 **235**, 147–156.
- 661 Klump J. V, Fitzgerald S.A. & Waples J.T. (2009) Benthic biogeochemical cycling, nutrient  
662 stoichiometry, and carbon and nitrogen mass balances in a eutrophic freshwater bay.  
663 *Limnology and Oceanography*, **54**, 692–712.
- 664 Kolattukudy P., Babel W. & Steinbüchel A. (2001) Polyesters in Higher Plants Biopolyesters.  
665 pp. 1–49. Springer Berlin / Heidelberg.
- 666 Koski-Vahala J. & Hartikainen H. (2001) Assessment of the risk of phosphorus loading due to  
667 resuspended sediment. *Journal of Environmental Quality*, **30**, 960–996.
- 668 Kosten S., Roland F., Marques D., Van Nes E. H., Mazzeo N., Sternberg L. D. L., Cole J. J.  
669 (2010) Climate-dependent CO<sub>2</sub> emissions from lakes. *Global Biogeochemical Cycles*,  
670 **24**, 7.
- 671 Kritzberg E.S., Langenheder S. & Lindstrom E.S. (2006) Influence of dissolved organic  
672 matter source on lake bacterioplankton structure and function - implications for seasonal  
673 dynamics of community composition. *FEMS Microbiology Ecology*, **56**, 406–417.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

674 Lacroix G., Boët P., Garnier J., Lescher-moutoué F., Pourriot R., & Testard P. (1989). Factors  
675 controlling the planktonic community in the shallow lake of Creteil, France.  
676 *Internationale Revue Der Gesamten Hydrobiologie*, **74(4)**, 353–370.

677 Lacroix G. & Lescher-Moutoué F. (1995). Spatial patterns of planktonic microcrustaceans in  
678 a small shallow lake. *Hydrobiologia*, **300-301**, 205–217.

679 Lowry O.H., Rosebrough N.J., Farr A.L. & Randall R.J. (1951) Protein measurement with the  
680 folin phenol reagent. *Journal of Biological Chemistry*, **193**, 265–275.

681 Martinova M. (1993) Nitrogen and phosphor compounds in bottom sediments: mechanisms of  
682 accumulation, transformation and release. *Hydrobiologia*, **252**, 1–22.

683 Masclaux H., Bec A., Kainz M.J., Desvillettes C., Jouve L. & Bourdier G. (2009) Combined  
684 effects of food quality and temperature on somatic growth and reproduction of two  
685 freshwater cladocerans. *Limnology and Oceanography*, **54**, 1323–1332.

686 Meckler A.N., Schubert C.J., Coxie G.L., Peiffer S. & Dittrich M. (2004) New organic matter  
687 degradation proxies: Valid in lake systems? *Limnology and Oceanography*, **49**, 2023–  
688 2033.

689 Mermillod-Blondin F., Gaudet J.-P., Gérino M., Desrosiers G. & Creuzé des Châtelliers M.  
690 (2003) Influence of macroinvertebrates on physico-chemical and microbial processes in  
691 hyporheic sediments. *Hydrological Processes*, **17**, 779–794.

692 Meyers P.A. (1997) Organic geochemical proxies of paleoceanographic, paleolimnologic, and  
693 paleoclimatic processes. *Organic Geochemistry*, **27**, 213–250.

- 694 Mortimer C. H. (1941) The exchange of dissolved substances between mud and water in  
695 lakes. *Journal of Ecology*, **29**, 280–329
- 696 Müller-Navarra D.C., Brett M.T., Liston A.M. & Goldman C.R. (2000) A highly unsaturated  
697 fatty acid predicts carbon transfer between primary producers and consumers. *Nature*,  
698 **403**, 74–77.
- 699 Nogaro G., Mermillod-Blondin F., Montuelle B., Boisson J.-C. & Gibert J. (2008)  
700 Chironomid larvae stimulate biogeochemical and microbial processes in a riverbed  
701 covered with fine sediment. *Aquatic Sciences*, **70**, 156–168.
- 702 Nürnberg G. K. (2009) Assessing internal phosphorus load – Problems to be solved. *Lake and*  
703 *Reservoir Management*, **25(4)**, 419–432.
- 704 Pettersson K. (1998) Mechanisms for internal loading of phosphorus in lakes. *Hydrobiologia*,  
705 **374**, 21–25.
- 706 Reynolds C.S. (1996) Phosphorus recycling in lakes: evidence from large limnetic enclosures  
707 for the importance of shallow sediments. *Freshwater Biology*, **35**, 623–645.
- 708 Sanford P.L. (1992) New sedimentation, resuspension, and burial. *Limnology and*  
709 *Oceanography*, **37**, 1164–1178.
- 710 Sarnelle O. (1999) Zooplankton effects on vertical particulate flux: Testable models and  
711 experimental results. *Limnology and Oceanography*, **44**, 357–370.
- 712 Schindler D.E., Carpenter S.R., Cole J.J., Kitchell J.F. & Pace M.L. (1997) Influence of food  
713 web structure on carbon exchange between lakes and the atmosphere. *Science*, **277**, 248–  
714 251.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

715 Schindler D.W. (1998) Replication versus realism: The need for ecosystem-scale  
716 experiments. *Ecosystems*, **1**, 323–334.

717 Schubert C.J., Niggemann J., Klockgether G. & Ferdelman T.G. (2005) Chlorin Index: A new  
718 parameter for organic matter freshness in sediments. *Geochemistry, Geophysics,*  
719 *Geosystems*, **6**.

720 Simon M., Grossart H.P., Schweitzer B. & Ploug H. (2002) Microbial ecology of organic  
721 aggregates in aquatic ecosystems. *Aquatic Microbial Ecology*, **28**, 175–211.

722 Sommer U. (1984) Sedimentation of principal phytoplankton species in Lake Constance.  
723 *Journal of Plankton Research*, **6**, 1–14.

724 Søndergaard M. (1988) Seasonal-variations in the loosely sorbed phosphorus fraction of the  
725 sediment of a shallow and hypereutrophic lake. *Environmental Geology and Water*  
726 *Sciences*, **11**, 115–121.

727 Søndergaard M., Jensen J.P. & Jeppesen E. (2003) Role of sediment and internal loading of  
728 phosphorus in shallow lakes. *Hydrobiologia*, **506-509**, 135–145.

729 Søndergaard M., Jeppesen E., Lauridsen T.L., Skov C., Van Nes E.H., Roijackers R., *et al.*  
730 (2007) Lake restoration: successes, failures and long-term effects. *Journal of Applied*  
731 *Ecology*, **44**, 1095–1105.

732 Spivak A.C., Canuel E.A., Duffy J.E. & Richardson J.P. (2007) Top-down and bottom-up  
733 controls on sediment organic matter composition in an experimental seagrass ecosystem.  
734 *Limnology and Oceanography*, **52**, 2595–2607.

- 735 Spivak A.C., Vanni M.J. & Mette E.M. (2011) Moving on up: can results from simple aquatic  
736 mesocosm experiments be applied across broad spatial scales? *Freshwater Biology*, **56**,  
737 279–291.
- 738 Urabe J., Kyle M., Makino W., Yoshida T., Andersen T. & Elser J.J. (2002) Reduced light  
739 increases herbivore production due to stoichiometric effects of light/nutrient balance.  
740 *Ecology*, **83**, 619–627.
- 741 Volkman J.K., Burton H.R., Everitt D.A. & Allen D.I. (1988) Pigment and lipid compositions  
742 of algal and bacterial communities in Ace Lake, Vestfold Hills, Antarctica.  
743 *Hydrobiologia*, **165**, 41–57.
- 744 Wakeham S. & Canuel E. (2006) Degradation and preservation of organic matter in marine  
745 sediments. In: *Marine Organic Matter: Biomarkers, Isotopes and DNA* (Eds. J.K. Volkman),  
746 pp. 295–321. The Handbook of Environmental Chemistry. Springer, Berlin/Heidelberg.
- 747 Weiss M. & Simon M. (1999) Consumption of labile dissolved organic matter by limnetic  
748 bacterioplankton: the relative significance of amino acids and carbohydrates. *Aquatic*  
749 *Microbial Ecology*, **17**, 1–12.
- 750 Yokoyama H., Takashi T., Ishihi Y. & Abo K. (2009) Effects of restricted feeding on growth  
751 of red sea bream and sedimentation of aquaculture wastes. *Aquaculture*, **286**, 80–88.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 1:** Selection criteria for the choice of the initial sediments. Elemental and biochemical composition (n = 1)

	C <sup>a</sup>	OC <sup>a</sup>	N <sup>a</sup>	OC/N	Sugars <sup>a</sup>	Proteins <sup>a</sup>	PUFAs <sup>b</sup>
S <sub>1</sub>	5.7	0.9	0.1	9.0	0.1	0.1	4.9
S <sub>2</sub>	23.1	9.7	2.4	4.0	0.7	1.5	12.6

<sup>a</sup> % of dry weight  
<sup>b</sup> % of total fatty acids.

**Table 2** Nutrient ( $\mu\text{g/L}$ ) concentrations of water sampled in May and June 2010 and DOC ( $\text{mg/L}$ ) concentrations of water sampled in May. Biomass ( $\text{mg L}^{-1}$ ), elemental and biochemical compositions (% of dry weight) of seston, zooplankton (sampled from February to June 2010). Sedimentation rates ( $\text{g DW m}^{-2} \text{d}^{-1}$ ), elemental and biochemical compositions (% of dry weight) of recent sediments (sampled in May and June 2010). Significant effects are in bold. Tukey's post-hoc tests were performed when the effect of sediment treatment was significant with the LME test.

	Means $\pm$ SD			Statistics							
	Treatment			Sediment						Time	Sediment $\times$ Time
				LME			Tukey (P values)				
	S <sub>0</sub>	S <sub>1</sub>	S <sub>2</sub>	P values	DF <sup>a</sup>	F values	S <sub>0</sub> -S <sub>1</sub>	S <sub>0</sub> -S <sub>2</sub>	S <sub>1</sub> -S <sub>2</sub>	P values	P values
<b>Water</b>											
Ammonium	28.5 $\pm$ 22.7	22.9 $\pm$ 22.1	32.7 $\pm$ 22.7	0.39	2, 21	0.98	-	-	-	<b>0.001</b>	0.52
Nitrates	4.6 $\pm$ 3.8	4.1 $\pm$ 2.2	4.5 $\pm$ 3.5	0.90	2, 21	0.10	-	-	-	<b>&lt; 0.001</b>	0.34
Phosphates	3.1 $\pm$ 5.3	3.8 $\pm$ 5.5	11.3 $\pm$ 10.3	<b>0.01</b>	2, 21	5.21	0.96	<b>0.01</b>	<b>0.02</b>	0.35	0.62
IN/P <sup>b</sup>	16 $\pm$ 21	18 $\pm$ 21	5 $\pm$ 5	0.24	2, 21	1.52	-	-	-	<b>&lt; 0.01</b>	0.35
DOC	4.6 $\pm$ 0.9	4.3 $\pm$ 0.9	5.8 $\pm$ 1.2	<b>0.02</b>	2	5.01	0.80	0.07	<b>0.02</b>	-	-
<b>Seston</b>											
Biomass	1.18 $\pm$ 0.63	1.40 $\pm$ 1.03	2.14 $\pm$ 1.49	<b>&lt; 0.005</b>	2, 21	6.92	0.57	<b>0.001</b>	<b>0.03</b>	<b>&lt; 0.0001</b>	0.11
C	30.3 $\pm$ 8.6	29.3 $\pm$ 8.8	32.2 $\pm$ 8.6	0.27	2, 21	1.39	-	-	-	<b>&lt; 0.001</b>	0.76
N	4.6 $\pm$ 1.9	4.7 $\pm$ 1.4	5.5 $\pm$ 1.5	<b>0.03</b>	2, 21	4.12	0.89	<b>0.02</b>	0.07	0.80	0.18
C/N ratio	6.6 $\pm$ 1.1	6.2 $\pm$ 1.0	5.9 $\pm$ 1.3	<b>&lt; 0.005</b>	2, 21	8.06	<b>0.03</b>	<b>&lt; 0.0001</b>	0.12	<b>&lt; 0.0001</b>	<b>&lt; 0.0001</b>
<b>Zooplankton</b>											
Biomass	0.27 $\pm$ 0.16	0.29 $\pm$ 0.19	0.51 $\pm$ 0.34	<b>0.002</b>	2, 21	8.87	0.99	<b>&lt; 0.0001</b>	<b>&lt; 0.0001</b>	<b>&lt; 0.0001</b>	<b>0.002</b>

C	42.8 ± 8.8	42.7 ± 4.6	44.3 ± 4.2	0.52	2, 21	0.67	-	-	-	< 0.01	0.84
N	7.9 ± 2.6	8.9 ± 2.0	8.0 ± 1.7	0.66	2, 21	0.43	-	-	-	< 0.0001	0.23
C/N ratio	5.4 ± 2.1	5.3 ± 1.4	5.3 ± 1.2	0.94	2, 21	0.06	-	-	-	< 0.0001	0.06
<i>Recent sediment</i>											
Sedimentation rate	0.77 ± 0.40	1.42 ± 0.96	1.17 ± 0.53	0.13	2, 21	2.25	-	-	-	0.09	0.65
C	24.8 ± 2.7	15.5 ± 3.7	26.9 ± 2.6	0.0001	2, 21	14.22	< 0.001	0.43	< 0.0001	0.03	0.10
N	2.8 ± 0.7	1.5 ± 0.5	3.3 ± 0.4	< 0.0001	2, 21	22.61	< 0.005	0.12	< 0.0001	0.97	0.65
C/N ratio	9.1 ± 2.4	10.3 ± 1.8	8.0 ± 1.1	< 0.005	2, 21	8.41	0.04	0.37	< 0.001	0.17	0.95
OC <sup>c</sup>	23.9 ± 2.6	14.4 ± 5.8	27.3 ± 2.2	0.02	2	8.93	0.05	0.58	0.01	-	-
OC/N <sup>c</sup>	8.9 ± 1.1	9.6 ± 5.8	8.0 ± 0.8	0.94	2	0.06	-	-	-	-	-
Sugars <sup>c</sup>	3.3 ± 0.3	2.7 ± 1.2	3.2 ± 0.8	0.36	2	1.22	-	-	-	-	-
Proteins <sup>c</sup>	2.2 ± 0.3	1.8 ± 0.6	3.2 ± 0.9	0.78	2	0.25	-	-	-	-	-

<sup>a</sup> Degrees of Freedom

<sup>b</sup> Inorganic Nitrogen (sum of ammonium and nitrates)/Phosphates ratio

<sup>c</sup> analyses were performed only on sediments sampled in May 2010

**Table 3** Mean lipid composition of seston, zooplankton and recent sediment sampled in May 2010. Statistical analyses were performed on the mean of the three sediment treatments for each compartment (mean  $\pm$  SD; n = 9 for seston, zooplankton and recent sediment). Bold values indicate significant differences between seston, zooplankton and/or recent sediment. Total was expressed as % of total lipids. Sub-classes of fatty acids and sterols were expressed as % of total fatty acids and sterols, respectively.

Mean	FAs					Sterols				OHS	OH-FAs	CHLOs
	Total	SAFA	MUFA	PUFA	BACTFA	Total	C <sub>27</sub> $\Delta^5$	C <sub>29</sub> $\Delta^5$	Stanols	Total	Total	Total
Seston	<b>75.4 <math>\pm</math> 9.3</b>	<b>74.2 <math>\pm</math> 5.4</b>	<b>11.9 <math>\pm</math> 4.4</b>	<b>4.2 <math>\pm</math> 2.8</b>	<b>9.7 <math>\pm</math> 1.5</b>	5.2 $\pm$ 2.0	<b>23.5 <math>\pm</math> 10.8</b>	<b>57.2 <math>\pm</math> 16.0</b>	<b>0.9 <math>\pm</math> 0.6</b>	<b>14.5 <math>\pm</math> 6.9</b>	<b>4.2 <math>\pm</math> 1.2</b>	<b>0.5 <math>\pm</math> 0.1</b>
Zooplankton	<b>87.3 <math>\pm</math> 7.0</b>	<b>52.8 <math>\pm</math> 5.3</b>	<b>21.5 <math>\pm</math> 5.3</b>	<b>19.4 <math>\pm</math> 4.9</b>	<b>6.4 <math>\pm</math> 1.2</b>	6.0 $\pm$ 4.6	<b>68.1 <math>\pm</math> 11.0</b>	<b>21.5 <math>\pm</math> 11.0</b>	<b>1.2 <math>\pm</math> 0.8</b>	<b>3.9 <math>\pm</math> 2.9</b>	<b>1.3 <math>\pm</math> 0.6</b>	<b>1.3 <math>\pm</math> 0.8</b>
Recent sediment	<b>75.0 <math>\pm</math> 7.7</b>	<b>56.2 <math>\pm</math> 3.2</b>	<b>24.0 <math>\pm</math> 2.2</b>	<b>11.1 <math>\pm</math> 2.5</b>	<b>8.7 <math>\pm</math> 1.9</b>	6.8 $\pm$ 4.1	<b>34.1 <math>\pm</math> 7.1</b>	<b>43.9 <math>\pm</math> 9.5</b>	<b>2.3 <math>\pm</math> 0.9</b>	<b>10.5 <math>\pm</math> 3.7</b>	<b>4.6 <math>\pm</math> 1.1</b>	<b>2.8 <math>\pm</math> 1.7</b>

FAs: fatty acids, SAFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acid, BACTFA: bacterial fatty acid =  $\Sigma$  15 :0 +  $\Sigma$  17 :0 + branched-chain FAs, OHS: *n*-alkanols, OH-FAs: sum of  $\alpha$ -,  $\beta$ - and  $\omega$ -hydroxy acids, CHLOs: chlorophyll-derived compounds.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 4** Mean specific composition of zooplankton communities (Individual L<sup>-1</sup>) sampled in February and May 2010. Mean ± SD. For each treatment and each date, counting was only performed on triplicates. Significant effects are in bold.

	Means ± SD			Statistics							
	Treatment			Sediment						Time	Sediment × Time
				LME			Tukey (P values)			P values	P values
	S <sub>0</sub>	S <sub>1</sub>	S <sub>2</sub>	P values	DF <sup>a</sup>	F values	S <sub>0</sub> -S <sub>1</sub>	S <sub>0</sub> -S <sub>2</sub>	S <sub>1</sub> -S <sub>2</sub>		
<b>Rotifers</b>	2.8 ± 2.8	1.7 ± 1.2	7.2 ± 15.4	0.86	2, 6	0.15	-	-	-	0.23	<b>0.002</b>
<b>Cladocerans</b>											
<i>Daphnia</i>	23.4 ± 22.2	11.4 ± 13.7	3.4 ± 4.2	0.11	2, 6	3.27	-	-	-	<b>0.066</b>	<b>0.24</b>
<i>Ceriodaphnia</i>	1.1 ± 2.2	0.1 ± 0.3	0.0 ± 0.1	0.18	2, 6	2.33	-	-	-	<b>0.0008</b>	<b>0.0054</b>
<i>Bosminidae</i>	1.3 ± 1.7	2.2 ± 3.9	2.2 ± 4.4	0.95	2, 6	0.05	-	-	-	< <b>0.0001</b>	0.94
Chydoridae	5.6 ± 6.2	2.8 ± 3.7	16.5 ± 23.3	<b>0.016</b>	2, 6	8.87	<b>0.76</b>	<b>0.02</b>	< <b>0.005</b>	< <b>0.0001</b>	< <b>0.0001</b>
<b>Copepods</b>											
Cyclopids	13.0 ± 14.0	7.5 ± 11.0	12.1 ± 12.5	0.33	2, 6	1.34	-	-	-	< <b>0.0001</b>	0.67
Calanoids	16.6 ± 10.8	10.3 ± 16.5	15.7 ± 18.6	0.41	2, 6	1.04	-	-	-	<b>0.0001</b>	<b>0.0008</b>
Nauplii	111.9 ± 111.5	61.0 ± 65.1	99.2 ± 124..0	0.11	2, 6	2.22	-	-	-	0.066	0.24

<sup>a</sup> Degrees of Freedom

1 Figure captions:

2

3 Figure 1: Seasonal variations of seston (a) and zooplankton (b) biomass and sedimentation  
4 rates (c) (mean  $\pm$  SE). White, grey and black bars represent the control treatment ( $S_0$ ),  $S_1$  and  
5  $S_2$  treatments respectively. Significant effects of time are represented by different letters.

6

7 Figure 2: Seasonal variations of carbon (a) and nitrogen (b) contents and C:N ratio (c) of the  
8 seston (mean  $\pm$  SE). White, grey and black bars represent the control treatment ( $S_0$ ),  $S_1$  and  $S_2$   
9 treatments respectively. Significant effects of time are represented by different letters.

10


11 Figure 3: Seasonal variations of carbon (a) and nitrogen (b) contents and C:N ratio (c) of the  
12 zooplankton (mean  $\pm$  SE). White, grey and black bars represent the control treatment ( $S_0$ ),  $S_1$   
13 and  $S_2$  treatment respectively. Significant effects of time are represented by different letters.

14

15 Figure 4: Variations of carbon (a) and nitrogen (b) contents and C:N ratio (c) of the short-term  
16 sediment (mean  $\pm$  SE). White, grey and black bars represent the control treatment ( $S_0$ ),  $S_1$  and  
17  $S_2$  treatment respectively. Significant effects of time are represented by different letters.

18

Figure 1



22 Figure 2






Figure 3



28 Figure 4



29

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Supplementary Table 1** Lipid composition of the initial sediments. Total was expressed as % of total lipids. Sub-classes of fatty acids, sterols, alkanols and hydroxy acids were expressed as % of total fatty acids, sterols, alkanols and hydroxy acids, respectively.

	FAs					Sterols				OHs			OH-FAs			CHLOs
	Total	SAFA	MUFA	PUFA	BACTFA	Total	$\Sigma\Delta^5$	$\Sigma\Delta^7$	stanols	Total	SC	LC	Total	$\alpha + \beta$	$\omega$	Total
S <sub>1</sub>	36.4	74.2	13.9	4.9	6.6	16.0	61.2	ND	38.8	26.3	5.1	94.9	8.5	71.4	26.9	4.5
S <sub>2</sub>	48.2	64.9	17.6	12.6	3.9	19.0	30.1	24.6	28.2	13.4	1.7	98.3	2.4	100.0	ND	9.8

FAs: fatty acids, SAFA: saturated fatty acid, MUFA: monounsaturated fatty acid, PUFA: polyunsaturated fatty acid, BACTFA: bacterial fatty acid =  $\Sigma 15:0 + \Sigma 17:0 +$  branched-chain FAs,  $\Sigma\Delta^5$ : sum of  $\Delta^5$ -sterols,  $\Sigma\Delta^7$ : sum of  $\Delta^7$ -sterols, OHs: alkanols, SC: short-chain alkanols (carbon number  $\leq 18$ ), LC: long-chain alkanols (carbon number  $\geq 20$ ), OH-FAs: hydroxy acids,  $\alpha + \beta$ : sum of  $\alpha$ - and  $\beta$ -hydroxy acids,  $\omega$ :  $\omega$ -hydroxy acids, CHLOs: chlorophyll-derived compounds, ND: non detected.

The quality (potential biodegradability) of the initial sediments was estimated using their contents in the constituents which were preferentially degraded by bacteria ie., proteins, sugars and PUFAs. The degradation state of these sediments was estimated using their contents in compounds related to the presence of bacteria ie. BACTFAs (Wakeham & Beier, 1991; Budge & Parrish, 1998) or to the bacterial activity ie.  $\alpha$ - and  $\beta$ -OH-FAs (Cranwell, 1981), and stanols, the saturated homologues of sterols (Gaskell & Eglinton, 1975).

The high relative abundance of  $\Delta^7$ -sterols in S<sub>2</sub> is consistent with the high contribution of Chlorophyceae to S<sub>2</sub> observed previously (Danger *et al.*, 2012). Indeed, these sterols are abundant in many Chlorophyceae (Volkman, 1986), and have been proposed as indicators of these algae (Cranwell, 1982). In contrast,  $\Delta^7$ -sterols were not detected in S<sub>1</sub>. 24-methyl- and 24-ethyl-cholest-5-enols, present in high amounts in S<sub>1</sub>, are abundant in algae and in vascular plants as well. So, the sterol distribution of S<sub>1</sub> cannot allow to discriminate between autochthonous and allochthonous inputs. The higher relative abundances of chlorophyll-derived compounds in S<sub>2</sub> than in S<sub>1</sub> corroborate also a higher phytoplanktonic contribution to S<sub>2</sub>. Indeed, such chlorophyll-derived compounds are usually considered as arising from the biodegradation of the phytol side-chain of chlorophyll (Rontani & Volkman, 2003).

The higher relative amounts of BACTFAs in S<sub>1</sub> than in S<sub>2</sub> suggest that S<sub>1</sub> was in a more advanced degradation state than S<sub>2</sub>. This is supported by higher amounts of  $\alpha$ - and  $\beta$ -OH-FAs and stanols in S<sub>1</sub> than in S<sub>2</sub>. Indeed,  $\alpha$ - and  $\beta$ -OH-FAs are usually related to microbial input in sediment (Cranwell, 1981), and stanols, the saturated homologues of sterols, are usually considered as formed through in situ microbial reduction of sterols (Gaskell & Eglinton, 1975).

**Supplementary Table 2** Fatty acid composition (% of total fatty acids) of the initial sediments S<sub>1</sub> and S<sub>2</sub> (n = 1).

SAFA	S <sub>1</sub>	S <sub>2</sub>	MUFA	S <sub>1</sub>	S <sub>2</sub>	PUFA	S <sub>1</sub>	S <sub>2</sub>	BACT	S <sub>1</sub>	S <sub>2</sub>
14:0	1.3	0.8	16:1 $\omega$ x	tr	tr	16:3 $\omega$ 3	ND	0.4	14:0 br	0.0	0.2
16:0	12.4	10.7	16:1 $\omega$ 7	5.1	1.7	16:2 $\omega$ x	ND	0.9	15:0 <sup>a</sup>	2.9	2.0
18:0	8.4	3.2	18:1 $\omega$ 9	4.8	9.5	18:3 $\omega$ 6	0.0	0.3	16:0 br	0.4	0.4
20:0	3.2	1.0	18:1 $\omega$ 7	3.0	2.8	18:4 $\omega$ 3	0.0	0.3	17:0 <sup>a</sup>	3.2	1.3
21:0	0.7	0.2	18:1 $\omega$ x	0.0	0.7	18:2 $\omega$ 6 (LIN)	1.0	1.6			
22:0	5.4	6.1	20:1 $\omega$ 9	0.9	1.2	18:3 $\omega$ 3 (ALA)	1.6	7.8			
23:0	1.8	0.5	22:1 $\omega$ x	0.0	0.2	20:4 $\omega$ 6 (ARA)	0.9	0.3			
24:0	12.6	9.7	24:1 $\omega$ x	0.0	0.5	20:5 $\omega$ 3 (EPA)	1.4	0.9			
25:0	1.6	0.7	26:1 $\omega$ x	0.0	0.8	22:5 $\omega$ 6	tr	0.1			
26:0	10.7	17.1				22:6 $\omega$ 3 (DHA)	tr	0.1			
27:0	1.2	0.9									
29:0	0.5	0.0									
30:0	2.9	1.5									
Subtotal	74.2	64.9		13.9	17.6		4.9	12.6		6.6	3.9
$\Sigma$ short-chain	22.1	14.7				$\Sigma$ short-chain	2.6	11.2			
$\Sigma$ long-chain	52.1	50.2				$\Sigma$ long-chain	2.3	1.4			



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

x: unknown position of double bounds. br: branched-chain FA. tr: traces, LIN: linoleic acid, ALA:  $\alpha$ -linolenic acid, ARA: arachidonic acid, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid. <sup>a</sup> Sum of branched and linear FAs.

Copy for Review

**Supplementary Table 3** Fatty acid composition (% of total FAs) of seston, zooplankton and recent sediment sampled in May 2010 (mean  $\pm$  SD; n = 3).

FA	Seston			Zooplankton			Recent sediment		
	S <sub>0</sub>	S <sub>1</sub>	S <sub>2</sub>	S <sub>0</sub>	S <sub>1</sub>	S <sub>2</sub>	S <sub>0</sub>	S <sub>1</sub>	S <sub>2</sub>
<b>SAFA</b>									
12:0	0.4 $\pm$ 0.2	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.0	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.2
14:0	4.1 $\pm$ 1.8	5.3 $\pm$ 2.5	3.2 $\pm$ 0.4	6.0 $\pm$ 5.1	7.2 $\pm$ 5.3	3.6 $\pm$ 1.4	2.4 $\pm$ 0.6	3.1 $\pm$ 1.5	3.0 $\pm$ 1.7
16:0	41.6 $\pm$ 3.3	40.6 $\pm$ 3.6	40.7 $\pm$ 2.7	32.1 $\pm$ 1.3	34.6 $\pm$ 2.0	36.7 $\pm$ 2.5	35.6 $\pm$ 1.7	33.5 $\pm$ 4.0	34.0 $\pm$ 2.3
18:0	17.5 $\pm$ 0.4	17.1 $\pm$ 4.0	19.3 $\pm$ 2.5	8.7 $\pm$ 0.7	11.3 $\pm$ 1.3	9.3 $\pm$ 0.8	11.6 $\pm$ 1.2	12.0 $\pm$ 0.9	9.6 $\pm$ 1.6
20:0	1.6 $\pm$ 0.1	1.6 $\pm$ 0.4	1.9 $\pm$ 0.2	0.5 $\pm$ 0.1	0.8 $\pm$ 0.2	0.5 $\pm$ 0.1	0.9 $\pm$ 0.8	1.6 $\pm$ 0.1	1.0 $\pm$ 0.2
21:0	0.5 $\pm$ 0.1	0.6 $\pm$ 0.2	0.6 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.0	0.2 $\pm$ 0.2	0.3 $\pm$ 0.0	0.2 $\pm$ 0.1
22:0	3.1 $\pm$ 0.0	3.6 $\pm$ 1.8	4.0 $\pm$ 0.2	0.8 $\pm$ 0.3	1.4 $\pm$ 0.5	0.7 $\pm$ 0.1	1.5 $\pm$ 1.5	3.2 $\pm$ 1.1	1.6 $\pm$ 1.1
23:0	0.7 $\pm$ 0.0	0.9 $\pm$ 0.5	0.9 $\pm$ 0.1	0.2 $\pm$ 0.0	0.3 $\pm$ 0.1	0.1 $\pm$ 0.0	0.3 $\pm$ 0.3	0.4 $\pm$ 0.2	0.3 $\pm$ 0.2
24:0	1.2 $\pm$ 0.2	1.5 $\pm$ 1.0	1.6 $\pm$ 0.3	0.3 $\pm$ 0.1	0.5 $\pm$ 0.4	0.2 $\pm$ 0.1	0.9 $\pm$ 0.9	1.2 $\pm$ 0.6	0.8 $\pm$ 0.4
25:0	0.3 $\pm$ 0.1	0.5 $\pm$ 0.4	0.5 $\pm$ 0.2	0.1 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.0	0.2 $\pm$ 0.2	0.2 $\pm$ 0.1	0.1 $\pm$ 0.1
26:0	0.3 $\pm$ 0.2	0.4 $\pm$ 0.4	0.4 $\pm$ 0.2	0.1 $\pm$ 0.1	0.2 $\pm$ 0.2	0.1 $\pm$ 0.1	0.5 $\pm$ 0.7	0.6 $\pm$ 0.5	0.5 $\pm$ 0.4
27:0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.0 $\pm$ 0.0	0.1 $\pm$ 0.0	0.0 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1
28:0	0.5 $\pm$ 0.4	0.7 $\pm$ 0.8	0.9 $\pm$ 0.7	0.1 $\pm$ 0.1	0.3 $\pm$ 0.3	0.1 $\pm$ 0.1	1.2 $\pm$ 1.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.4
30:0	0.5 $\pm$ 0.6	0.7 $\pm$ 0.9	0.7 $\pm$ 0.8	0.1 $\pm$ 0.1	0.3 $\pm$ 0.4	0.1 $\pm$ 0.1	1.3 $\pm$ 1.4	0.5 $\pm$ 0.8	1.8 $\pm$ 3.0
32:0	0.2 $\pm$ 0.1	0.2 $\pm$ 0.2	0.3 $\pm$ 0.3	0.0 $\pm$ 0.0	0.1 $\pm$ 0.1	0.0 $\pm$ 0.0	0.1 $\pm$ 0.1	0.2 $\pm$ 0.2	0.1 $\pm$ 0.1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Subtotal	72.6 ± 3.8	74.3 ± 9.5	75.6 ± 2.5	49.3 ± 2.8	57.3 ± 7.0	51.8 ± 2.4	57.2 ± 2.0	57.5 ± 2.2	54.1 ± 4.5
Σshort-chain	63.6 ± 5.0	63.4 ± 3.8	63.7 ± 5.0	47.1 ± 3.3	53.2 ± 8.5	49.8 ± 3.1	49.9 ± 2.8	48.8 ± 6.4	47.1 ± 3.9
Σlong-chain	9.0 ± 1.6	10.9 ± 6.6	12.0 ± 2.7	2.2 ± 0.8	4.1 ± 2.2	2.0 ± 0.7	7.3 ± 4.4	8.8 ± 4.2	7.0 ± 5.8
MUFA									
15:1ωx	0.5 ± 0.2	0.4 ± 0.2	0.3 ± 0.2	0.2 ± 0.1	0.4 ± 0.3	0.5 ± 0.1	0.2 ± 0.1	0.4 ± 0.1	0.6 ± 0.5
16:1ω7	3.6 ± 1.3	4.1 ± 2.9	3.3 ± 0.8	4.1 ± 0.6	3.9 ± 1.5	5.2 ± 2.4	4.4 ± 0.3	4.8 ± 1.9	5.1 ± 1.7
16:1ωx	0.2 ± 0.3	0.1 ± 0.1	0.3 ± 0.2	0.4 ± 0.1	1.0 ± 0.5	1.2 ± 0.4	1.5 ± 0.3	1.5 ± 0.6	2.5 ± 0.5
17:1ωx	0.3 ± 0.5	0.1 ± 0.1	0.0 ± 0.0	0.4 ± 0.1	0.3 ± 0.2	0.4 ± 0.3	0.6 ± 0.2	0.6 ± 0.2	1.2 ± 0.6
18:1ω9	6.4 ± 1.6	2.3 ± 1.6	5.2 ± 0.7	10.0 ± 2.8	9.8 ± 2.1	9.4 ± 2.0	8.1 ± 0.9	10.6 ± 0.9	8.5 ± 0.7
18:1ω7	2.2 ± 2.0	0.9 ± 0.7	1.5 ± 1.3	5.2 ± 1.5	4.0 ± 2.4	6.2 ± 2.5	6.5 ± 0.8	4.3 ± 1.3	4.2 ± 0.6
18:1ωx	0.3 ± 0.4	1.7 ± 2.8	0.6 ± 1.1	0.1 ± 0.0	0.1 ± 0.1	0.1 ± 0.1	0.4 ± 0.1	0.4 ± 0.1	2.9 ± 1.8
20:1ωx	0.2 ± 0.1	0.3 ± 0.2	0.2 ± 0.0	0.4 ± 0.0	0.3 ± 0.0	0.3 ± 0.0	0.3 ± 0.0	0.5 ± 0.3	0.4 ± 0.1
20:1ω9	0.1 ± 0.1	0.0 ± 0.1	0.2 ± 0.0	0.1 ± 0.0	0.0 ± 0.0	0.1 ± 0.0	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.0
22:1ωx	0.1 ± 0.2	0.2 ± 0.4	0.0 ± 0.1	0.0 ± 0.1	0.0 ± 0.1	0.0 ± 0.1	0.7 ± 0.9	0.3 ± 0.3	0.1 ± 0.1
Subtotal	13.9 ± 3.2	10.2 ± 7.3	11.8 ± 1.8	21.1 ± 4.4	19.9 ± 5.7	23.5 ± 7.0	22.9 ± 1.5	23.5 ± 0.4	25.6 ± 3.2
PUFA									
16:xω3	0.1 ± 0.1	0.2 ± 0.2	ND	0.2 ± 0.1	0.1 ± 0.0	0.2 ± 0.1	0.4 ± 0.1	0.3 ± 0.1	0.3 ± 0.0
16:3ω3	0.1 ± 0.1	0.1 ± 0.1	ND	0.3 ± 0.1	0.2 ± 0.1	0.5 ± 0.0	0.3 ± 0.0	0.4 ± 0.2	0.7 ± 0.2
16:2ωx	0.0 ± 0.1	0.1 ± 0.1	ND	0.1 ± 0.0	0.2 ± 0.0	0.1 ± 0.1	ND	ND	ND
18:3ω6	ND	ND	ND	0.3 ± 0.1	0.1 ± 0.0	0.2 ± 0.1	0.2 ± 0.0	0.2 ± 0.1	0.1 ± 0.0

18:2 $\omega$ 3	0.5 $\pm$ 0.2	0.3 $\pm$ 0.2	0.1 $\pm$ 0.2	1.9 $\pm$ 0.6	1.0 $\pm$ 0.4	1.5 $\pm$ 1.1	0.8 $\pm$ 0.2	0.5 $\pm$ 0.2	0.4 $\pm$ 0.1
18:2 $\omega$ 6 (LIN)	1.5 $\pm$ 1.0	0.9 $\pm$ 0.6	0.8 $\pm$ 0.2	6.1 $\pm$ 1.7	5.4 $\pm$ 1.0	2.9 $\pm$ 0.7	5.7 $\pm$ 0.9	4.6 $\pm$ 1.5	3.4 $\pm$ 0.2
18:3 $\omega$ 3 (ALA)	1.7 $\pm$ 1.2	2.9 $\pm$ 3.5	1.4 $\pm$ 0.8	7.6 $\pm$ 2.5	4.6 $\pm$ 1.2	6.7 $\pm$ 1.7	3.5 $\pm$ 0.4	3.7 $\pm$ 1.8	4.1 $\pm$ 1.5
18:2 $\omega$ x	0.3 $\pm$ 0.1	0.8 $\pm$ 0.5	0.7 $\pm$ 0.4	1.1 $\pm$ 1.0	0.4 $\pm$ 0.3	0.7 $\pm$ 0.2	0.4 $\pm$ 0.1	0.4 $\pm$ 0.2	0.4 $\pm$ 0.1
20:4 $\omega$ 6 (ARA)	0.1 $\pm$ 0.1	ND	0.0 $\pm$ 0.1	1.4 $\pm$ 0.3	1.1 $\pm$ 0.1	0.6 $\pm$ 0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1
20:5 $\omega$ 3 (EPA)	0.0 $\pm$ 0.1	ND	0.0 $\pm$ 0.0	2.4 $\pm$ 0.7	1.6 $\pm$ 0.6	2.1 $\pm$ 1.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.1	0.3 $\pm$ 0.1
20:2 $\omega$ x	ND	ND	ND	0.6 $\pm$ 0.1	0.4 $\pm$ 0.1	0.2 $\pm$ 0.1	ND	ND	0.1 $\pm$ 0.1
20:3 $\omega$ 3	ND	ND	ND	0.2 $\pm$ 0.1	0.1 $\pm$ 0.1	0.3 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1
20:3 $\omega$ x	ND	ND	ND	0.2 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.0 $\pm$ 0.0	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1
22:5 $\omega$ 6	ND	ND	ND	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0
22:6 $\omega$ 3 (DHA)	ND	ND	ND	1.3 $\pm$ 0.4	0.8 $\pm$ 0.5	1.0 $\pm$ 0.7	0.1 $\pm$ 0.1	0.0 $\pm$ 0.1	0.1 $\pm$ 0.0
22:5 $\omega$ 3	ND	ND	ND	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	0.1 $\pm$ 0.0	ND	ND	ND
Subtotal	4.2 $\pm$ 2.2	5.3 $\pm$ 4.7	3.1 $\pm$ 1.1	24.1 $\pm$ 3.1	16.6 $\pm$ 2.0	17.3 $\pm$ 5.5	12.0 $\pm$ 1.4	10.9 $\pm$ 4.1	10.2 $\pm$ 1.6
$\Sigma$ short-chain	4.1 $\pm$ 2.0	5.3 $\pm$ 4.7	3.0 $\pm$ 1.0	17.6 $\pm$ 2.3	12.1 $\pm$ 0.9	12.7 $\pm$ 3.3	11.3 $\pm$ 1.5	10.0 $\pm$ 3.8	9.3 $\pm$ 1.9
$\Sigma$ long-chain	0.1 $\pm$ 0.2	0.0 $\pm$ 0.0	0.1 $\pm$ 0.1	6.5 $\pm$ 0.9	4.6 $\pm$ 1.3	4.5 $\pm$ 2.2	0.8 $\pm$ 0.1	0.9 $\pm$ 0.3	0.9 $\pm$ 0.3
<b>BACTFA</b>									
14:0 br	0.2 $\pm$ 0.1	0.1 $\pm$ 0.0	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.0	0.2 $\pm$ 0.1	0.3 $\pm$ 0.1	0.3 $\pm$ 0.1	0.3 $\pm$ 0.2
15:0 br	1.9 $\pm$ 0.6	2.3 $\pm$ 1.4	1.8 $\pm$ 0.2	1.8 $\pm$ 0.2	1.5 $\pm$ 0.2	2.2 $\pm$ 0.3	2.0 $\pm$ 0.3	2.3 $\pm$ 1.3	4.0 $\pm$ 1.8
15:0	2.0 $\pm$ 0.6	2.0 $\pm$ 0.1	1.9 $\pm$ 0.2	0.9 $\pm$ 0.1	1.3 $\pm$ 0.7	1.2 $\pm$ 0.2	1.4 $\pm$ 0.3	1.5 $\pm$ 0.4	1.9 $\pm$ 0.6
16:0 br	0.2 $\pm$ 0.0	0.2 $\pm$ 0.2	0.2 $\pm$ 0.2	0.3 $\pm$ 0.1	0.2 $\pm$ 0.2	0.3 $\pm$ 0.1	0.5 $\pm$ 0.0	0.3 $\pm$ 0.1	0.5 $\pm$ 0.1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

17:0 br	1.0 ± 0.2	1.1 ± 0.4	1.0 ± 0.4	0.7 ± 0.1	0.8 ± 0.3	1.2 ± 0.7	1.5 ± 0.9	1.2 ± 0.0	1.2 ± 0.2
17:0	3.4 ± 0.1	4.2 ± 0.4	3.8 ± 0.1	1.6 ± 0.1	2.0 ± 0.7	2.1 ± 0.7	2.1 ± 1.0	2.1 ± 0.2	1.8 ± 0.0
19:0	0.4 ± 0.2	0.3 ± 0.0	0.4 ± 0.1	0.1 ± 0.0	0.1 ± 0.1	0.1 ± 0.0	0.2 ± 0.0	0.3 ± 0.2	0.3 ± 0.1
Subtotal	9.3 ± 1.2	10.3 ± 2.7	9.5 ± 0.1	5.5 ± 0.1	6.1 ± 1.5	7.5 ± 1.0	7.9 ± 1.1	8.1 ± 1.7	10.1 ± 2.4

ND non detected. x: unknown position of double bounds. br: branched-chain FA

**Supplementary References:**

- Budge S.M. & Parrish C.C. (1998) Lipid biogeochemistry of plankton, settling matter and sediments in Trinity Bay, Newfoundland. II. Fatty acids. *Organic Geochemistry*, **29**, 1547–1559.
- Cranwell P.A. (1981) Diagenesis of free and bound lipids in terrestrial detritus deposited in a lacustrine sediment. *Organic Geochemistry*, **3**, 79–89.
- Cranwell P.A. (1982) Lipids of aquatic sediments and sedimenting particulates. *Progress in Lipid Research*, **21**, 271–308.
- Danger M., Allard B., Arnous M., Carrias J.-F., Mériquet J., Ten-Hage L., *et al.* (2012) Effects of food-web structure on the quantity and the elemental quality of sedimenting material in shallow lakes. *Hydrobiologia*, **679**, 251–266.
- Gaskell S.J. & Eglinton G. (1975) Rapid hydrogenation of sterols in a contemporary lacustrine sediment. *Nature*, **254**, 209–211.
- Rontani J.-F. & Volkman J.K. (2003) Phytol degradation products as biogeochemical tracers in aquatic environments. *Organic Geochemistry*, **34**, 1–35.
- Volkman J.K. (1986) A review of sterol markers for marine and terrigenous organic matter. *Organic Geochemistry*, **9**, 83–99.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Wakeham S. G., & Beier J. A. (1991). Fatty-acids and sterols biomarkers as indicator of particulate matter source and alteration processes in the Black-sea. *Deep-Sea Research Part a-Oceanographic Research Papers*, **38**, S943–S968.

Copy for Review